

AD-A084 775

PROFESSIONAL STAFF ASSOCIATION OF LAC/USC MEDICAL CEN--ETC F/G 6/15
STUDIES ON THE BIOLOGY, VENOM AND ULTRASTRUCTURE OF SELECTED VE--ETC(U)
APR 80 F E RUSSELL

N00014-78-C-0206

NL

UNCLASSIFIED

AD
A084 775



END
DATE
FILMED
6-80
DTIC



PROFESSIONAL STAFF ASSOCIATION Los Angeles County/ University of Southern California Medical Center

LEVEL II

(9)

Final Technical

Report to

OCEANIC BIOLOGY PROGRAM

OFFICE OF NAVAL RESEARCH

ADA 084775

Submitted by

(10)

Findlay E. Russell

Director, Laboratory of Neurological Research
Los Angeles County/University of Southern California
Medical Center,

Through the

Professional Staff Association

6

STUDIES ON THE BIOLOGY, VENOM AND ULTRASTRUCTURE OF SELECTED
VENOMOUS FISHES, INCLUDING THE SCORPIONFISHES AND STINGRAYS.

OFFICE OF NAVAL RESEARCH
CONTRACT / N00014-78-C-0206

(15)

MAY 27 1980

D

C

DDC FILE COPY

25 Apr 1980

(12) 7

Findlay E. Russell, M.D., Ph.D.
Principal Investigator

13
100014-78-C-0206
100014-78-C-0206

80 5 19 111

110679

mt

STUDIES ON THE BIOLOGY, VENOM AND ULTRASTRUCTURE OF SELECTED
VENOMOUS FISHES, INCLUDING THE SCORPIONFISHES AND STINGRAYS

Findlay E. Russell
Director, Laboratory of Neurological Research
Los Angeles County-University of Southern California
Medical Center, Los Angeles, California

Work Unit No. NR 104-793

Contract N00014-78-C-0206

During the past two years the principal research under the contract has been directed toward several problems:

1) A continued study of the ultrastructure of the venom apparatus of the stingrays Urobatis halleri and Dasyatis sabina, and the scorpionfish Scorpaena guttata; 2) determination of certain physiopharmacologic changes provoked by the venom of the scorpionfishes; and 3) the histology and ultrastructure of skin from the soapfishes, Grammistidae.

With respect to current studies on the ultrastructure of the stingray venom apparatus, these have confirmed our earlier observations that the subunit distribution in stingray venom cell "microtubules" varies considerably within a single vacuole, in marked contrast to the tubulin polymers described as "conventional cytoplasmic microtubules."

The improvement in the methods we have developed aided in the definition of these structures in thin sections and permits a more precise estimate of their diameter. While diameters are variable, they are substantially less than estimates for other intracellular microtubules. The chemistry of these microtubules has yet to be determined but by their anomalous location and fine structure it is suggested that they are polymers of non-tubulin protein(s). One protein component (150,000 daltons) was identified in Urobatis halleri venom. The variability in subunit

distribution in venom cell microtubules may indicate that a number of proteins are sequestered and polymerized within the vacuoles.

The improvements in the techniques we have developed for stabilizing the venoms of the scorpionfishes has permitted us to do a number of physiopharmacologic studies that heretofore have not been possible. Essentially, the procedure is as follows: Using the separation method previously developed in our laboratory, venom was extracted in 0.05 M sodium phosphate buffer, pH 7.4, and 10^{-3} M Cleland's reagent concentrated two-fold by ultrafiltration, using a UM-2 (Amicon) membrane under 75 psi N_2 pressure.

A cardiovascular survey model designed to study hypotension was employed to measure arterial and venous pressures; respirations; and blood pH, PO_2 , PCO_2 and hematocrit; and total serum protein. The results indicated a fairly consistent pattern and has led to the employ of more definitive isolated-tissue segment preparations. This work is currently being completed. There is a close relationship between our findings and those we have observed in patients stung by the scorpionfishes.

Studies on several species of soapfishes (Grammistidae) indicated that there is a general pattern of three secretory cells in the epidermis of these fishes, as distinguished by histochemical staining and ultrastructural appearance. One secretory cell is a large, oval, PAS negative, Sudan black (SB) negative, homogenous, glassy appearing cell extending from the deep epithelial layers to the skin surface. Viewed with the electron microscope its contents appear as highly packed tiny granules. The second secretory cell is round and contains tightly packed large, PAS positive, SB negative granules, which empty through narrow openings. This cell is in the surface epithelium and is thought to be the mucus secreting cells found in all teleost skin. The third type

of secreting cell is the elongated cell extending from the deep epithelium to the surface, where it empties its irregular granules which are PAS negative and SB positive. It appears that the fish's toxin is probably a product of this third cell type with synergism with the second type.

The following papers, supported by the contract were published or are in press:

Chao, R.L.C., Parker, J.W., Russell, F.E. and Hashimota, Y. (1978)
"Ultrastructure of the skin of the soapfish Grammistes sexlineatus."
In, Toxins: Animal, Plant and Microbial. Rosenberg, P. (Ed.) p. 449-508, Pergamon Press, Oxford.

Smith, D.S., Cayer, M. and Russell, F.E. (1978)
"Fine structure of stingray epidermis with special reference to a unique microtubular component of the venom secreting cells." In, Toxins: Animal, Plant and Microbial. Rosenberg, P. (Ed.) p. 565-582, Pergamon Press: Oxford.

Russell, F.E. (1978)
Hazardous marine life. Part I. Venomous marine animals. Hyperbaric & Undersea Medicine 1: 1-8.

Russell, F.E., Smith, D.S., Cayer, M. and Gonzalez, H.C. (1978)
Behavior, ecology and toxicity of venomous marine fishes. ONR Prog. Rept. ACR-228, p. 91 December (Abst.).

Maretic, Z., Russell, F.E. and Ladavac, J. (1979)
Epidemic of stings by the jellyfish Pelagia noctiluca in the Adriatic.

Abstract of the 6th International Symposium on Animal, Plant and Microbial Toxins, Uppsala, August. Toxicon 17, 115.

Russell, F.E. (1979)

Epidemiologija I Terapije Uboda I Ujeda Otrovnih Zivotinja U. Sad.
II Kongres Udruzenja Toksikologa Jugoslavije, Portoroz, Jugoslavije,
p. 27.

Russell, F.E., Smith, D.S., Cayer, M. and Gonzalez, H.C. (1979)

Behavior, Ecology and Toxicity of Venomous Marine Fishes. ONR Prog.
Rept. ACR-230, p. 99 December. (Abst.)

Russell, F.E. (1980)

The marine organism sting mystery. (Questions and Answers) J.A.M.A.
243: 1573.

Coats, J.A., Pattabhiraman, T.R., Russell, F.E. and Gonzalez, H.C. (1980)

Some physiopharmacologic properties of scorpionfish venom. Proc. west.
Pharmacol. Soc. (In Press)

Sloan, C. and Russell, F.E. (1980)

Ultrastructure of the venom apparatus of the scorpionfish Scorpaena
guttata. Toxicon. (In Press)

Smith, D.S., Cayer, M. and Russell, F.E. (1980)

Tannic acid staining of "microtubules" in venom-secreting cells of the
stingray. Toxicon. (In Press)

Dymsza, H., Shimizu, Y., Russell, F.E. and Graham, H.D. (1980)

"Poisonous Marine Animals." In, The Safety of Foods. Graham, H.D., (Ed.)
p. 625-651, AVI: Westport, Connecticut.

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER 1 - Final	2. GOVT ACCESSION NO. AD-A084 775	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) Studies on the biology, venom and ultrastructure of selected venomous fishes including the scorpionfishes and stingrays		5. TYPE OF REPORT & PERIOD COVERED Final - Technical
7. AUTHOR(s) Findlay E. Ruaaell ^{SS} , M.S., Ph.D.		6. CONTRACT OR GRANT NUMBER(s) N00014-78-C-0206
9. PERFORMING ORGANIZATION NAME AND ADDRESS Professional Staff Association of the LAC/USC Medical Center 1739 Griffin Ave., Los Angeles, CA. 90031		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS
11. CONTROLLING OFFICE NAME AND ADDRESS		12. REPORT DATE April 25, 1980
		13. NUMBER OF PAGES 2
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) Commanding Officer Office of Naval Research Branch Office 1030 East Green Street Pasadena, CA. 91101		15. SECURITY CLASS. (of this report) Unclassified
16. DISTRIBUTION STATEMENT (of this Report) Unlimited		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number)		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) During the past two years the principal research under the contract has been directed toward several problems: (1) A continued study of the ultrastructure of the venom apparatus of the stingrays <i>Urobatis halleri</i> and <i>Dasyatis sabina</i> , and the scorpionfish <i>Scorpaena Guttata</i> , (2) Determination of certain physiopharmacologic changes provoked by the venom of the scorpionfishes,		

DD FORM 1473
1 JAN 73EDITION OF 1 NOV 68 IS OBSOLETE
S/N 0102-014-6601

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

80 5 19 111

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

(3) The histology and ultrastructure of skin from the soapfishes, Grammistidae. With respect to current studies on the ultrastructure of the stingray venom apparatus, these have confirmed our earlier observations that the subunit distribution in stingray venom cell microtubules varies considerably within a single vacuole, in marked contrast to the tubulin polymers described as conventional cytoplasmic microtubules. The improvement in the methods we have developed, aided in the definition of these structures in thin sections and permits a more precise estimate of their diameter. While diameters are variable, they are substantially less than estimates for other intracellular microtubules. The chemistry of these microtubules has yet to be determined but by their anomalous location and fine structure it is suggested that they are polymers of non-tubulin protein(s). One protein component (150,000 daltons) was identified in Urobatis halleri venom. The variability in subunit distribution in venom cell microtubules may indicate that a number of proteins are sequestered and polymerized within the vacuoles. The improvements in the techniques we have developed for stabilizing the venoms of the scorpionfishes has permitted us to do a number of physiopharmacologic studies that heretofore have not been possible. Essentially, the procedure is as follows: Using the separation method previously developed in our laboratory, venom was extracted in 0.05 M sodium phosphate buffer, pH 7.4, and 10-3 M Cleland's reagent concentrated two-fold by ultrafiltration, using a UM-2 (Amicon) membrane under 75 psi N2 pressure. A cardiovascular survey model designed to study hypotension was employed to measure arterial and venous pressures/respirations; and blood pH, PO2, PCO2 and hematocrit; and total serum protein. The results indicated a fairly consistent pattern and has led to the employ of more definitive isolated-tissue segment preparations. This work is currently being completed. There is a close relationship between our findings and those we have observed in patients stung by the scorpionfishes.

Accession For	
NRIS	GRAMI
INDS	CAB
Unannounced	
Classification	
by	
Distribution	
Availability	
Dist	Availability
A	Special